

# Propalin®

## Short Response Time, High Success Rate

Propalin® is a veterinary only product that directly targets the urethral muscle and increases the bladder's ability to retain urine. Within a week, you should notice significant improvement in your dog's condition. As urethral muscle tone increases over the next few weeks, progress should continue to be maintained over time.

- **Proven Effective and Safe** – In clinical trials side effects of Propalin® were reported as being infrequent, mild, and temporary<sup>1</sup>
- 89.7% Overall Positive Response Rate
- Fast Results You Can See
- The Only Phenylpropanolamine with Documented Efficacy

**(Note:** If your dog's symptoms have not improved within a few weeks on the medication, you should consult your veterinarian.)



(Available in 25 mg, 50 mg,  
and 75 mg chewable tablets)

<sup>1</sup>Leddy, M.C., Weaver, Jr., and Eyett-Burton, C.  
Long-term study of phenylpropanolamine in the control of urinary incontinence in the bitch

Propalin® A Solution to the Problem.

The worldwide treatment of choice for urinary incontinence in dogs



- ☑ Easy to administer
- ☑ Well tolerated
- ☑ Non-hormonal
- ☑ Proven safe and effective for long-term use
- ☑ Low incidence of side effects
- ☑ Rapid improvement
- ☑ High success rate



Vétoquinol USA Inc.  
101 Lincoln Avenue  
Buena, NJ 08310  
1-800-267-5707  
[www.propalinusa.com](http://www.propalinusa.com)



## Important Information About Your Pet's Urinary Incontinence and **HOW** Propalin® Can Help.

Propalin® Ahead of the Crowd, Accept no Substitutes.

The worldwide treatment of choice for urinary incontinence in dogs



 Vétoquinol  
*a Sign of Passion*

# Propalin® (phenylpropanolamine)

## Safely and Effectively Manage Urinary Incontinence...and Preserve the Pet-Owner Bond

### Understanding urinary incontinence


If you find “wet spots” on bedding or furnishings where your dog has been sleeping or lying down, or if its hindquarters are wet, it could be a sign of urinary incontinence.

What causes urinary incontinence?

While urinary incontinence can indicate an underlying condition or infection, diagnostic tests don't always provide a clear answer.

Often, the best way to confirm urinary incontinence is by trial treatment. If symptoms are alleviated, the cause is most likely what is known as sphincter incompetence, a condition that affects your dog's ability to control urine.

In many cases, a decrease in certain hormones due to the natural aging process may be a cause.



### Is your dog at risk?

Although urinary incontinence can affect dogs of any age and of both sexes, it seems to occur more frequently in the following:

- ☑ Dogs 6 years of age and older
- ☑ Spayed females
- ☑ Mid to large-size breeds
- ☑ Overweight dogs
- ☑ Dogs affected by underlying conditions

**Decreased urethral muscle tone**



Loss of urethral muscle tone can result in urinary incontinence.

**Normal urethral muscle tone**



Cross section of the urethra

The urethral muscle acts as a valve to prevent urine from escaping the bladder.

### A new lease on life

Sphincter incompetence is not a disease. It does not cause pain and it is manageable. Help your dog regain bladder control and restore harmony in your home with easy to administer Propalin®.

### A few accident prevention tips

- Confine your dog to an area that can be easily cleaned, such as the laundry room, basement or garage—especially when it is left alone for extended periods.
- Cover your dog's favorite sleeping spots with old blankets and/or towels that can be washed easily. Place a waterproof pad under the blankets or towels as extra “insurance”.
- You may want to consider using special disposable pads and/or “diapers”, depending on the situation.

And above all, remember that it's not your dog's fault...and that things will soon be better!



**Propalin®** Remember the Yellow Pack, Not the Puddle.

The worldwide treatment of choice for urinary incontinence in dogs